

Anaphylaxis Emergency Action Plan

Patient Name: _____ Age: _____

Allergies: _____

Asthma Yes (*high risk for severe reaction*) No

Additional health problems besides anaphylaxis: _____

Concurrent medications: _____

	Symptoms of Anaphylaxis
MOUTH	itching, swelling of lips and/or tongue
THROAT*	itching, tightness/closure, hoarseness
SKIN	itching, hives, redness, swelling
GUT	vomiting, diarrhea, cramps
LUNG*	shortness of breath, cough, wheeze
HEART*	weak pulse, dizziness, passing out

*Only a few symptoms may be present. Severity of symptoms can change quickly.
Some symptoms can be life-threatening. ACT FAST!

Emergency Action Steps - DO NOT HESITATE TO GIVE EPINEPHRINE!

1. Inject epinephrine in thigh using (check one): Adrenaclick (0.3 mg) Adrenaclick (0.15 mg)
 EpiPen Jr (0.15 mg) EpiPen (0.3 mg)
 Twinject (0.15 mg) Twinject (0.3 mg)

Other medication/dose/route: _____

IMPORTANT: ASTHMA INHALERS AND/OR ANTIHISTAMINES CAN'T BE DEPENDED ON IN ANAPHYLAXIS.

2. Call 911 or rescue squad (before calling contact)

3. Emergency contact #1: home _____ work _____ cell _____

Emergency contact #2: home _____ work _____ cell _____

Emergency contact #3: home _____ work _____ cell _____

Comments: _____

() The student has been instructed by me in the proper use of his/her medication and may carry and self-administer the above medications while on school property or at school related events.
 () The student should NOT be allowed to carry and self-administer his/her medications. (Recommended for all elementary students.)

 Doctor's Signature/Date/Phone Number

 Parent's Signature (for individuals under age 18 yrs)/Date